Lateral manipulation for the positioning of molecular guests within the confinements of a highly stable self-assembled organic surface network

Stöhr, Meike; Wahl, Markus; Spillmann, Hannes; Gade, Lutz H.; Jung, Thomas A.

Published in:
Small

DOI:
10.1002/smll.200700099

Publication date:
2007

Supporting Information

Supporting Information

Lateral manipulation for the positioning of molecular guests within the Confinements of a Highly Stable Self-Assembled Organic Surface Network

by

Meike Stöhr,*[a] Markus Wahl,[a] Hannes Spillmann,[a] Lutz H. Gade*[b]
and Thomas A. Jung*[c]

[a] NCCR Nanoscale Science and Institute of Physics, University of Basel, Klingelbergstr. 82, 4056 Basel, Switzerland. E-mail: meike.stoehr@unibas.ch

[b] Anorganisch-Chemisches Institut, Universität Heidelberg, Im Neuenheimer Feld 270, 69120 Heidelberg, Germany. E-mail: lutz.gade@uni-hd.de

[c] Laboratory for Micro- and Nanostructures, Paul-Scherrer-Institute, 5232 Villigen, Switzerland. E-mail: thomas.jung@psi.ch

---

---

1) STM-Images of C_{60} deposited onto dehydro-DPDI on Cu(111) recorded at ambient temperature

Figure S1



2004-11-12_31, RT, 50 nm x 33 nm

Figure S2



2005-03-16_21, RT, 50 nm x 50 nm

2) STM-Images of C_{60} deposited onto dehydro-DPDI on Cu(111) recorded at 77K

Figure S3



2005-07-27_07, 77 K, 100 nm x 100 nm

Figure S4



2005-07-28_20, 77 K, 58 nm x 58 nm

3) STM-Images of C_{60} deposited at high density onto dehydro-DPDI on Cu(111) recorded at 77K

Figure S5



2005-09028_19, 77 K, 28nm x 28nm



2005-09-28_43, 77 K, 10nm x 10nm

Figure S6



2006-01-23_13, 77 K, 30nm x 30nm